

CAROLINE CALDERONE BAISLEY
DIRECTOR OF HEALTH



MICHAEL LONG, Director
DIVISION OF ENVIRONMENTAL SERVICES
DOUGLAS SERAFIN, Laboratory
DEBORAH TRAVERS, Director
DIVISION OF FAMILY HEALTH
THOMAS MAHONEY, Director
SPECIAL CLINICAL SERVICES
STEPHANIE PAULMENO, Director
COMMUNITY HEALTH PLANNING

DEPARTMENT OF HEALTH

Swine Flu: FAQ's

QUESTIONS AND ANSWERS:

Swine Influenza and You

Updated April 29, 2009

What is swine flu?

Swine Influenza (swine flu) is a respiratory disease of pigs caused by type A influenza viruses. Outbreaks of swine flu happen regularly in pigs. People do not normally get swine flu, but human infections can and do happen. Most commonly, human cases of swine flu happen in people who are around pigs but it's possible for swine flu viruses to spread from person to person also.

Are there human infections with swine flu in the U.S.?

In late March and early April 2009, cases of human infection with swine influenza A (H1N1) viruses were first reported in Southern California and near San Antonio, Texas. As of April 29th, 2009 there are 91 confirmed cases of human swine flu with one reported fatality in Texas in 9 states (California, Kansas, Ohio, Texas, New York, Massachusetts, Michigan, Nevada and Michigan.) There are two suspected, but not confirmed cases in Connecticut. CDC and local and state health agencies are working together to investigate this situation.

Is this swine flu virus contagious?

CDC has determined that this virus is contagious and is spreading from human to human. However, at this time, it not known how easily the virus spreads between people.

What are the signs and symptoms of swine flu in people?

The symptoms of swine flu in people are similar to the symptoms of regular human flu and include fever, cough, sore throat, body aches, headache, chills and fatigue. Some people have reported diarrhea and vomiting associated with swine flu. In the past, severe illness (pneumonia and respiratory failure) and deaths have been reported with swine flu infection in people. Like seasonal flu, swine flu may cause a worsening of underlying chronic medical conditions.

How serious is swine flu infection?

Like seasonal flu, swine flu in humans can vary in severity from mild to severe. Between 2005 until January 2009, 12 human cases of swine flu were detected in the U.S. with no deaths occurring. However, swine flu infection can be serious. In September 1988, a previously healthy 32-year-old pregnant woman in Wisconsin was hospitalized for pneumonia after being infected with swine flu and died 8 days later. A swine flu outbreak in Fort Dix, New Jersey occurred in 1976 that caused more than 200 cases with serious illness in several people and one death.

How do you catch swine flu?

Spread of swine flu can occur in two ways:

- Through contact with infected pigs or environments contaminated with swine flu viruses.
- Through contact with a person with swine flu. Human-to-human spread of swine flu has been documented also and is thought to occur in the same way as seasonal flu. Influenza is thought to spread mainly person-to-person through coughing or sneezing of infected people.

Are there medicines to treat swine flu?

Yes. CDC recommends the use of oseltamivir or zanamivir for the treatment and/or prevention of infection with these swine influenza viruses. Antiviral drugs are prescription medicines (pills, liquid or an inhaler) that fight against the flu by keeping flu viruses from reproducing in your body. If you get sick, antiviral drugs can make your illness milder and make you feel better faster. They may also prevent serious flu complications. Testing and treatment with antiviral drugs will be determined by your health care provider and work best if started soon after getting sick (within 2 days of symptoms).

How long can an infected person spread swine flu to others?

People with swine influenza virus infection should be considered potentially contagious as long as they are symptomatic and possible for up to 1 one prior to onset of illness and 7 days following illness onset. Children, especially younger children, might potentially be contagious for longer periods.

What can I do to protect myself from getting sick?

There is no vaccine available right now to protect against swine flu. There are everyday actions that can help prevent the spread of germs that cause respiratory illnesses like influenza. Take these everyday steps to protect your health:

- Cover your nose and mouth with a tissue when you cough or sneeze. Throw the tissue in the trash after you use it.
- Wash your hands often with soap and water, especially after you cough or sneeze. Alcohol-based hand cleaners are also effective.
- Try to avoid close contact with sick people.
- If you get sick with influenza, CDC recommends that you stay home from work or school and limit contact with others to keep from infecting them. Avoid touching your eyes, nose or mouth. Germs spread this way.

What should I do if I get sick?

If you live in San Diego County or Imperial County California or Guadalupe County, Texas and become ill with influenza-like symptoms, including fever, body aches, runny nose, sore throat, nausea, or vomiting or diarrhea, you may want to contact their health care provider, particularly if you are worried about your symptoms. Your health care provider will determine whether influenza testing or treatment is needed.

If you are sick, you should stay home and avoid contact with other people as much as possible to keep from spreading your illness to others.

If you become ill and experience any of the following warning signs, seek emergency medical care.

In children emergency warning signs that need urgent medical attention include:

- Fast breathing or trouble breathing
- Bluish skin color
- Not drinking enough fluids
- Not waking up or not interacting
- Being so irritable that the child does not want to be held
- Flu-like symptoms improve but then return with fever and worse cough
- Fever with a rash

In adults, emergency warning signs that need urgent medical attention include:

- Difficulty breathing or shortness of breath
- Pain or pressure in the chest or abdomen
- Sudden dizziness
- Confusion
- Severe or persistent vomiting

Can I get swine influenza from eating or preparing pork?

No. Swine influenza viruses are not spread by food. You cannot get swine influenza from eating pork or pork products. Eating properly handled and cooked pork products is safe.

How many swine flu viruses are there?

Like all influenza viruses, swine flu viruses change constantly. Pigs can be infected by avian influenza and human influenza viruses as well as swine influenza viruses. When influenza viruses from different species infect pigs, the viruses can reassort (i.e. swap genes) and new viruses that are a mix of swine, human and/or avian influenza viruses can emerge. Over the years, different variations of swine flu viruses have emerged. At this time, there are four main influenza type A virus subtypes that have been isolated in pigs: H1N1, H1N2, H3N2, and H3N1. However, most of the recently isolated influenza viruses from pigs have been H1N1 viruses.

How common is swine flu infection in humans?

In the past, CDC received reports of approximately one human swine influenza virus infection every one to two years in the U.S., but from December 2005 through February 2009, 12 cases of human infection with swine influenza have been reported.

How does swine flu spread?

Influenza viruses can be directly transmitted from pigs to people and from people to pigs. Human infection with flu viruses from pigs are most likely to occur when people are in close proximity to infected pigs, such as in pig barns and livestock exhibits housing pigs at fairs. Human-to-human transmission of swine flu can also occur. This is thought to occur in the same way as seasonal flu occurs in people, which is mainly person-to-person transmission through coughing or sneezing of people infected with the influenza virus. People may become infected by touching something with flu viruses on it and then touching their mouth or nose.

What other examples of swine flu outbreaks are there?

Probably the most well known is an outbreak of swine flu among soldiers in Fort Dix, New Jersey in 1976. The virus caused disease with x-ray evidence of pneumonia in at least 4 soldiers and 1 death; all of these patients had previously been healthy. The virus was transmitted to close contacts in a basic training environment, with limited transmission outside the basic training group. The virus is thought to have circulated for a month and disappeared. The source of the virus, the exact time of its introduction into Fort Dix, and factors limiting its spread and duration are unknown. The Fort Dix outbreak may have been caused by introduction of an animal virus into a stressed human population in close contact in crowded facilities during the winter. The swine influenza A virus collected from a Fort Dix soldier was named A/New Jersey/76 (Hsw1N1).

Is the H1N1 swine flu virus the same as human H1N1 viruses?

No. The H1N1 swine flu viruses are antigenically very different from human H1N1 viruses and, therefore, vaccines for human seasonal flu would not provide protection from H1N1 swine flu viruses.

How does swine flu spread among pigs?

Swine flu viruses are thought to be spread mostly through close contact among pigs and possibly from contaminated objects moving between infected and uninfected pigs. Herds with continuous swine flu infections and herds that are vaccinated against swine flu may have sporadic disease, or may show only mild or no symptoms of infection.

What are signs of swine flu in pigs?

Signs of swine flu in pigs can include sudden onset of fever, depression, coughing (barking), discharge from the nose or eyes, sneezing, breathing difficulties, eye redness or inflammation, and going off feed.

How common is swine flu among pigs?

H1N1 and H3N2 swine flu viruses are endemic among pig populations in the United States and something that the industry deals with routinely. Outbreaks among pigs normally occur in colder weather months (late fall and winter) and sometimes with the introduction of new pigs into susceptible herds. Studies have shown that the swine flu H1N1 is common throughout pig populations worldwide, with 25 percent of animals showing antibody evidence of infection. In the U.S. studies have shown that 30 percent of the pig population has antibody evidence of having had H1N1 infection. More specifically, 51 percent of pigs in the north-central U.S. have been shown to have antibody evidence of infection with swine H1N1. Human infections with swine flu H1N1 viruses are rare. There is currently no way to differentiate antibody produced in response to flu vaccination in pigs from antibody made in response to pig infections with swine H1N1 influenza.

While H1N1 swine viruses have been known to circulate among pig populations since at least 1930, H3N2 influenza viruses did not begin circulating among US pigs until 1998. The H3N2 viruses initially were introduced into the pig population from humans. The current swine flu H3N2 viruses are closely related to human H3N2 viruses.

Is there a vaccine for swine flu?

Vaccines are available to be given to pigs to prevent swine influenza. There is no vaccine to protect humans from swine flu. The seasonal influenza vaccine will likely help provide partial protection against swine H3N2, but not swine H1N1 viruses.

Related Links And Information:

Greenwich Department of Health Flu Checklist, www.greenwichct.org or call 203-622-7838

Connecticut Department of Public Health, www.ct.gov/dph or call Info Line at 211.

Greenwich Hospital, www.greenhosp.org/ or call 203-863-4455.

Center for Disease Control; and Prevention General information,

www.cdc.gov

www.cdc.gov/swineflu/general_info.htm

www.cdc.gov/flu/swine/index.htm

Center for Disease Control and Prevention Swine Flu Information,

www.cdc.gov/swineflu/casedef_swineflu.htm

www.cdc.gov/swineflu/guidelines_infection_control.htm

Center for Disease Control and Prevention Travel Tips,

www.cdc.gov/travel/

*Centers for Disease Control and Prevention 1600 Clifton Rd. Atlanta, GA 30333, USA
800-CDC-INFO (800-232-4636) TTY: (888) 232-6348, 24 Hours/Every Day - cdcinfo@cdc.gov*